

temperature conditions of formation of the assemblages are qualitatively consistent with that model. Significant aspects of that model may therefore now be considered verified by the evidence of the natural mineral assemblages discussed herein.

The primitive 'pyrolite' chemical composition of the mantle has been taken (somewhat arbitrarily) to be equivalent to a mixture of one part of basalt to three parts of dunite. The rocks discussed in this paper possess chemical compositions closely approaching that model. It should be emphasized, however, that any parent rock characterized by basalt-dunite ratios between 1:1 and 1:10 would probably be capable of crystallizing in each of the four mineral assemblages, according to the specific *P-T* conditions.

Accordingly, if ultrabasic rocks in this composition range do, in fact, predominate in the upper mantle, large-scale mineralogical zoning controlled by the *P-T* distribution will be inevitable. The upper mantle cannot, therefore, be regarded as a homogeneous region possessing a characteristic set of physical parameters (such as density and seismic velocity). Implications of this mineralogical zoning with respect to the presence or absence of a low-velocity zone, and to regional variations in seismic velocity profiles and surface heat flow, were outlined in previous papers [Ringwood, 1962a, b]. Further progress now awaits the direct experimental determination of the *P-T* fields of the four assemblages, as well as a better knowledge of the elastic properties of the individual minerals of these assemblages, as a function of temperature and pressure.

REFERENCES

- Boyd, F. R., and J. L. England, Minerals of the mantle, *Carnegie Inst. Wash. Year Book* 69, 47-55, 1960.
- Brown, G. M., The layered ultrabasic rocks of Rhum, Inner Hebrides, *Phil. Trans. Roy. Soc. London, B*, 240, 1-53, 1957.
- Carstens, C. W., Norske peridotiter I, II, *Norsk Geol. Tidsskr.*, 5, 1-73, 1920.
- Clark, S. P., and J. de Neufville, Crystalline phases in the system diopside-CaAl₂SiO₆-silica at high pressures (abstract), *J. Geophys. Res.*, 67, 3550, 1962.
- Davidson, C. F., The Archaean rocks of the Rodil district, South Harris, *Trans. Roy. Soc. Edin-*
- Dawson, J. B., Basutoland kimberlites, *Bull. Geol. Soc. Am.*, 73, 545-560, 1962.
- Du Rietz, T., Peridotites, serpentines, and soapstones of northern Sweden, *Geol. Foren. Forh.*, H 2, 133-260, 1935.
- Eskola, P., On the eclogites of Norway, *Vidensk. Skr. 1. Math.-Nat. Kl. Kristiania*, 14-88, 1921.
- Eskola, P., On the granulites of Lapland, *Am. J. Sci., Bowen Vol.*, 133-171, 1952.
- Gass, I. G., Ultrabasic pillow lavas from Cyprus, *Geol. Mag.*, 95, 241-251, 1958.
- Green, D. H., Petrogenesis of the peridotite at the Lizard, Cornwall, and The high-temperature aureole of the Lizard peridotite, Cornwall, papers in press from Ph.D. thesis, University of Cambridge, 1963.
- Green, D. H., Ultramafic breccias from Hie Mura valley, Eastern Papua, *Geol. Mag.*, 98, 1-25, 1961.
- Hess, H. H., A primary peridotite magma, *Am. J. Sci.*, 35, 321-344, 1938.
- Hess, H. H., Serpentine, orogeny and epeirogeny, in *Crust of the Earth*, *Geol. Soc. Am. Spec. Paper* 62, 1955.
- Hess, H. H., The Stillwater igneous complex, *Geol. Soc. Am. Mem.* 80, 1960a.
- Hess, H. H., Caribbean research project progress report, *Bull. Geol. Soc. Am.*, 71, 235-240, 1960b.
- Holmes, A., A contribution to the petrography of the kimberlites, *Trans. Geol. Soc. S. Africa*, 36, 379-428, 1936.
- Howie, R. H., The geochemistry of the charnockite series of Madras, India, *Trans. Roy. Soc. Edinburgh*, 62, 725-768, 1955.
- Johannesen, A., *A Descriptive Petrography of the Igneous Rocks*, vol. 4, University of Chicago Press, 1938.
- Nockolds, S. R., Average chemical compositions of some igneous rocks, *Bull. Geol. Soc. Am.*, 65, 1007-1032, 1954.
- Ringwood, A. E., On the chemical evolution and densities of the planets, *Geochim. Cosmochim. Acta*, 15, 257-283, 1959.
- Ringwood, A. E., A model for the upper mantle, *J. Geophys. Res.*, 67, 857-866, 1962a.
- Ringwood, A. E., A model for the upper mantle, 2, *J. Geophys. Res.*, 67, 4473-4477, 1962b.
- Reed, J. J., Chemical and modal composition of dunite from Dun Mt., Nelson, New Zealand, *J. Geol. Geophys.*, 2, 916-918, 1959.
- Robertson, E. C., F. Birch, and G. J. F. MacDonald, Experimental determination of jadeite stability relations to 25,000 bars, *Am. J. Sci.*, 255, 115, 1957.
- Ross, C. S., M. D. Foster, and A. T. Myers, Origin of dunites and of olivine-rich inclusions in basaltic rocks, *Am. Mineralogist*, 32, 693-737, 1954.
- Stanley, E. R., Lherzolite and olivine from Mt. Gambier, *Trans. Roy. Soc. S. Australia*, 34, 63-68, 1910.
- Stenvik, K. J., The properties of olivine and its uses for refractories and moulding sands, *Iron*