

by major rift or fault systems. More voluminous magmas, which are also usually products of much greater degrees of partial melting, form when there is diapiric upwelling of material from the low velocity zone to higher levels in the mantle or lower crust (Green and Ringwood 1967; Green 1970).

Magma types ranging from tholeiitic picrite to olivine melilitite and olivine trachybasalt are assigned to a petrogenetic grid in which the depth (pressure) of magma segregation, the degree of partial melting of the source rock, the water content, and approximate temperature of the magma, are specified. While this model can account for variations in major element abundances and normative mineralogy among basalts, there are variations in abundances of K, Rb, Ba, the rare-earths, etc. (incompatible elements), which are inconsistent with a model involving a constant source composition for all mantle derived basalts. Abundances of incompatible elements in the magmas are partly determined by the source composition at magma segregation but may be further enriched by wall-rock reaction processes in the upper mantle or in the continental crust. Source inhomogeneity appears to be required to account for variations in incompatible element abundances and such inhomogeneity may be a characteristic feature within the low velocity zone, the upper part of which is considered to have 'enriched' and the lower part a 'depleted' pattern of incompatible element contents.

REFERENCES

- BOYD, F. R., 1967. Electron probe study of diopsidic pyroxenes from kimberlite. *Carnegie Inst. Washington Year Bk.*, **65**, 252-260.
- BULTITUDE, R. J. and GREEN, D. H., 1968. Experimental studies at high pressures on the origin of olivine nephelinite and olivine melilitite nephelinitic magmas. *Earth Planetary Sci. Lett.*, **3**, 325-337.
- BULTITUDE, R. J. and GREEN, D. H., 1970. Experimental study of crystal-liquid relationship at high pressures in olivine nephelinite and basanite compositions. *J. Petrol.* (In press).
- CARSWELL, D. A. and DAWSON, J. B., 1970. Garnet peridotite xenoliths in South African kimberlite pipes and their petrogenesis. *Contr. Mineral. Petrol.*, **25**, 163-184.
- CLARK, S. P. and RINGWOOD, A. E., 1964. Density distribution and constitution of the mantle. *Rev. Geophys.*, **2**, 35-88.
- COOPER, J. A. and GREEN, D. H., 1969. Lead isotope measurements on lherzolite inclusions and host basanites from Western Victoria, Australia. *Earth Planetary Sci. Lett.*, **6**, 69-76.
- FREY, F. A., 1969. Rare earth abundances in a high temperature peridotite intrusion. *Geochim. Cosmochim. Acta.*, **33**, 1429-1447.
- GAST, P. W., 1968. Trace element fractionation and the origin of tholeiitic and alkaline magmas types. *Geochim. Cosmochim. Acta.*, **32**, 1,057-1,086.
- GREEN, D. H., 1969a. The origin of basaltic and nephelinitic magmas in the earth's mantle. *Tectonophysics*, **7**, 409-422.
- GREEN, D. H., 1969b. A review of experimental evidence on the origin of basaltic and nephelinitic magmas. *Phys. Earth Planetary Interiors*, **3**, 221-235.
- GREEN, D. H., 1970. Compositions of basaltic magmas as indicators of conditions of origin: application to oceanic volcanism. *Phil. Trans. Roy. Soc. London*, (In press).
- GREEN, D. H. and HIBBERSON, W., 1969. Experimental duplication of conditions of precipitation of high pressure phenocrysts in a basaltic magma. *Phys. Earth Planetary Interiors*, **3**, 247-254.
- GREEN, D. H. and HIBBERSON, W., 1970. Instability of plagioclase in peridotite at high pressures. *Lithos*, **3**, 209-221.
- GREEN, D. H. and RINGWOOD, A. E., 1967a. The stability fields of aluminous pyroxene peridotite and garnet peridotite and their relevance in upper mantle structure. *Earth Planetary Sci. Lett.*, **3**, 151-160.

- GREEN, D. H. and RINGWOOD, A. E., 1967b. The genesis of basaltic magmas. *Contr.Mineral.Petrol.*, **15**, 103 - 190.
- GREEN, D. H. and RINGWOOD, A. E., 1967c. An experimental study of the gabbro to eclogite transformation and its petrological applications. *Geochim. Cosmochim.Acta.*, **31**, 767 - 833.
- GREEN, D. H. and RINGWOOD, A. E., 1969. Mineralogy of peridotitic compositions under upper mantle conditions. *Phys.EarthPlanetaryInteriors*, **3**, 359 - 371.
- JACKSON, E. D. and WRIGHT, T. L., 1970. Xenoliths in the Honolulu Volcanic Series, Hawaii. *J.Petrol.*, **11**, 405 - 430.
- KUNO, H., 1960. High-alumina basalt. *J.Petrol.*, **1**, 121 - 145.
- KUSHIRO, I., 1969. Discussion of paper "The origin of basaltic and nephelinitic magmas in the earth's mantle". *Tectonophysics*, **7**, 427 - 436.
- KUSHIRO, I., YODER, H. S. and NISHIKAWA, M., 1968. Effect of water on the melting of enstatite. *Geol.Soc.Amer.Bull.*, **79**, 1,685 - 1,692.
- MACDONALD, G. A. and KATSURA, T., 1961. Variations in the lava of the 1959 eruption in Kilauea Iki. *Pacific Sci.*, **15**, 358 - 369.
- MURATA, K. J. and RICHTER, D. H., 1966. Chemistry of the lavas of the 1959 - 60 eruption of Kilauea volcano, Hawaii. *U.S.Geol.Surv.Profess.Papers*, **537-A**, 1 - 26.
- NOCKOLDS, S. R., 1954. Average chemical compositions of some igneous rocks. *Geol.Soc.Amer.Bull.*, **65**, 1,007 - 1,032.
- O'HARA, M. J., 1968. The bearing of phase equilibria studies in synthetic and natural systems on the origin and evolution of basic and ultrabasic rocks. *Earth Sci.Rev.* **4**, 69 - 133.
- O'HARA, M. J. and YODER, H. S., 1964. Formation and fractionation of basic magmas at high pressures. *Scottish J.Geol.*, **3**, 67 - 117.
- RINGWOOD, A. E., 1962. A model for the upper mantle. *J.Geophys.Res.*, **67**, 857 - 867.
- RINGWOOD, A. E., 1966a. The chemical composition and origin of the earth. p. 287 - 356 in *Advances in Earth Science*, ed. P. M. Hurley. M.I.T. Press, Cambridge, Mass.
- RINGWOOD, A. E., 1966b. The mineralogy of the mantle. p. 357 - 417 in *Advances in Earth Science*, ed. P. M. Hurley. M.I.T. Press, Cambridge, Mass.
- TILLEY, C. E., 1950. Some aspects of magmatic evolution. *Quart.J.Geol.Soc., London*, **106**, 37 - 61.

ACKNOWLEDGMENTS

The writer is grateful to Professor Sylvester-Bradley and Dr. M. J. Le Bas for their invitation to deliver this lecture and it is hoped that its publication will be of some value in summarizing a current view of a rapidly advancing and somewhat controversial aspect of igneous petrology. I also take this opportunity to gratefully acknowledge the stimulating collaboration with Professor A. E. Ringwood, the instruction and stimulus of Professor C. E. Tilley, and the essential role of Professor J. C. Jaeger and academic and technical colleagues at the Australian National University.