

## DISCUSSION

Bowen and Schairer [1935], using the Clapeyron-Clausius equation, calculated an initial slope for the melting curve of fayalite to be  $3.5^\circ/\text{kb}$ . The latent heat of fusion of fayalite used in their calculation was derived from the solidus and liquidus curves for the  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  system. The volume change of fayalite upon melting was roughly estimated by extrapolating the refractive index data of both crystal and glass in the  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  system. They also stated that direct measurement by Goranson [Bowen and Schairer, 1935, p. 208] showed the rise of melting point to be approximately twice as much as the calculated value. The slope of  $7.5^\circ/\text{kb}$  obtained in the present study essentially verified Goranson's earlier result. Inaccurate estimation of both the latent heat and the volume change of fayalite may be chiefly responsible for the disagreement between the calculated and experimental values.

Experimental results on the olivine-spinel transition in  $\text{Fe}_2\text{SiO}_4$  obtained in this study essentially agree with those by Boyd and England [1960], who used a piston-cylinder type of high-pressure apparatus with similar fur-

nace assembly. The stability field of  $\text{Fe}_2\text{SiO}_4$  spinel shown in Figure 2 consistently involves the experimental conditions of  $1400^\circ\text{C}$  at 65 kb which were determined by them to be in the spinel field.

The average slope of the melting curve for fayalite is  $5.1^\circ/\text{kb}$ , being close to that for forsterite,  $4.77^\circ/\text{kb}$ , in the pressure range up to about 50 kb [Davis and England, 1964]. It is expected, therefore, that the melting relations in the system  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  at pressures up to 62 kb would be essentially the same as those at atmospheric pressure. At pressures higher than 62 kb, however,  $(\text{Mg},\text{Fe})_2\text{SiO}_4$  spinel appears to be on the liquidus, and the solidus of olivine solid solution is intersected by the solvus between olivine and spinel solid solutions. Figure 3, the estimated equilibrium diagram of the system  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  at about 75 kb, shows such relations. With increasing pressure the solvus shifts toward  $\text{Mg}_2\text{SiO}_4$ , and the liquidus and solidus of spinel solid solution become wider.

The remarkable inflection in the melting curve of  $\text{Fe}_2\text{SiO}_4$ , which coincides with the melting of  $\text{Fe}_2\text{SiO}_4$  spinel, is of considerable geophysical importance. This suggests that a similar inflection should be anticipated in the melting-point gradient of the earth's mantle, provided that the olivine-spinel transformation takes place in the transition zone of the mantle. Positive support of the above inference is found in calculations on the melting curve of the mantle. Uffen [1952] estimated quite independently the change in melting temperature of the mantle with depth using seismic data on the basis of the Einstein-Debye theory of solids and Lindemann's theory of fusion. Clark [1963], considering thermodynamics and the effect of the transition zone, derived a melting curve for the mantle. We can find a considerable inflection in both melting curves around the transition zone. Since the chemical composition of the olivine in the mantle is generally supposed to be in the  $\text{Mg}_2\text{SiO}_4$ -rich side of the  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  system, it is highly desirable to extend this work to a comprehensive study of the fusion of the  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  system at high pressures.

*Acknowledgments.* We wish to express our thanks to Dr. T. Katsura and Mr. K. Kitayama of Tokyo Institute of Technology for their kind




Fig. 3. Equilibrium diagram of the system  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  at pressures near 75 kb. Melting point of forsterite is obtained by extrapolating the curve given by Davis and England [1964]. The liquidus and solidus are assumed. The solvus is based on the olivine-spinel transition in the system  $\text{Mg}_2\text{SiO}_4$ - $\text{Fe}_2\text{SiO}_4$  at 800, 1000, and  $1200^\circ\text{C}$  (Akimoto and Fujisawa, unpublished data).

help in preparing the fayalite sample used in this work. We are especially grateful to Dr. S. Aramaki of the Earthquake Research Institute, University of Tokyo, who offered helpful suggestions on petrology in the early stage of this work.

## REFERENCES

- Akimoto, S., and H. Fujisawa, Demonstration of the electrical conductivity jump produced by the olivine-spinel transition, *J. Geophys. Res.*, **70**, 443-449, 1965.
- Akimoto, S., H. Fujisawa, and T. Katsura, The olivine-spinel transition in  $\text{Fe}_2\text{SiO}_4$  and  $\text{Ni}_2\text{SiO}_4$ , *J. Geophys. Res.*, **70**, 1969-1977, 1965.
- Bowen, N. L., and J. F. Schairer, The system,  $\text{MgO-FeO-SiO}_2$ , *Am. J. Sci.*, **29**, 151-217, 1935.
- Boyd, F. R., and J. L. England, The fayalite- $\text{Fe}_2\text{SiO}_4$  spinel transition, *Carnegie Inst. Wash. Year Book 59*, pp. 48-49, 1960.
- Boyd, F. R., and J. L. England, Effect of pressure on the melting of diopside,  $\text{CaMgSi}_2\text{O}_6$ , and albite,  $\text{NaAlSi}_3\text{O}_8$ , in the range up to 50 kilobars, *J. Geophys. Res.*, **68**, 311-323, 1963.
- Clark, S. P., Jr., Variation of density in the earth and the melting curve in the mantle, in *The Earth Sciences*, edited by T. W. Donnelly, pp. 5-42, University of Chicago Press, 1963.
- Davis, B. T. C., and J. L. England, The melting of forsterite up to 50 kilobars, *J. Geophys. Res.*, **69**, 1113-1116, 1964.
- Uffen, R. J., A method of estimating the melting-point gradient in the earth's mantle, *Trans. Am. Geophys. Union*, **33**, 893-896, 1952.

(Received July 2, 1966;  
revised September 17, 1966.)